

Mimophis mahfalensis (common big-eyed snake) predation on *Furcifer major* (giant carpet chameleon) in Isalo, Madagascar

GONÇALO M. ROSA^{1,2,3*}, JOÃO CASTELO BRANCO³, HENRIQUE COUTO³,
PAOLO EUSEBIO BERGÒ⁴ & FRANCO ANDREONE⁴

¹Durrell Institute of Conservation and Ecology, School of Anthropology and Conservation, University of Kent, Canterbury CT2 7NR, Kent, UK

²Institute of Zoology, Zoological Society of London, Regent's Park, London NW1 4RY, UK

³Centro de Biologia Ambiental, Faculdade de Ciências da Universidade de Lisboa, Bloco C2, 1749-016 Lisbon, Portugal

⁴Museo Regionale di Scienze Naturali, Via G. Giolitti, 36, 10123 Turin, Italy

*corresponding author Email: goncalo.m.rosa@gmail.com

Snakes are carnivorous and many are specialist feeders on a particular type of prey. Some of the prey may include other reptiles such as lizards and even other snakes (Bauchot, 1994; Mercurio et al., 2006). In Madagascar there are about 100 species of non-marine colubrids in which the Lamprophiidae family represents over 80 of the described species (Cadle, 2003; Laduke et al., 2010; Crottini et al., 2012). *Mimophis mahfalensis* (Grandidier, 1867), is an endemic colubrid with a widespread distribution in Madagascar (except on the east coast) (Glaw & Vences, 2007). The diet of this snake includes frogs, other snakes and lizards (Glaw & Vences, 2007; Jenkins et al., 2009).

During our ongoing research on the herpetofauna of the Isalo Massif, central-southern Madagascar, an adult individual of *M. mahfalensis* was observed preying on an adult male of *Furcifer major* (Brygoo, 1971) on 14 February 2011. The episode took place in Malaso, 22°35'29.28"S; 45°21'26.82"E (Fianarantsoa Province, Antsohy Fivondronona, Ranohira Firaisana).

When the team approached the animals (at 13:00), the snake was already wrapped tightly around the chameleon, which was still alive. For few minutes the prey was moving but struggling in failed attempts (Fig. 1A). After the lizard apparently abandoned attempts at resistance and showed no signs of life, the snake adjusted its prey in order to start swallowing from the head (Fig. 1B). Twenty minutes later the snake began swallowing the lizard head, first briefly pausing when reaching the front limbs (Fig. 1C). The rest of the swallowing process took about 40 mins (Fig. 1D), which made the total observation last over an hour since the animals were first spotted.

F. major is a common chameleon endemic to Madagascar (Raxworthy et al., 2003; Florio et al., 2012). As part of *F. lateralis* complex, it has also been reported as a prey for other snake species (e.g. *Ithycyphus oursi*; Domergue, 1986; Crottini et al., 2010). Although a not recorded previously as a prey species for *M. mahfalensis*, it is known to prey on other chameleons (e.g. *F. labordi*; Andriamandimbarisoa, 2007). This opisthoglyphous snake (Rosa et al., 2014), uses a combination of mild envenomation and constriction to immobilise and kill its prey. As commented by Domergue (1989), *M. mahfalensis* seems to have a preference for saurians. Given how

common both species are, particularly in the Isalo region, predation on *F. major* by *M. mahfalensis* is likely a frequent event, especially when *F. major* moves through open areas.

ACKNOWLEDGEMENTS

We thank our guide Anicet for the help in the field. The study was carried out in collaboration with MNP and PBZT. We are grateful to the Malagasy authorities, in particular the Ministère de l'Environnement et des Eaux et Forêts, for research permits. Fieldwork was funded by Mohamed bin Zayed Species Conservation Fund, Gondwana Conservation and Research, EDGE, and Zoological Society of London.

REFERENCES

- Andriamandimbarisoa, L.N. (2007). Contribution à l'étude de l'histoire naturelle de trois espèces de caméléons de la région de Toliary, *Furcifer verrucosus* (Cuvier, 1829), *Furcifer labordi* (Grandidier, 1872) et *Furcifer antimena* (Grandidier, 1872): biologie, écologie et éthologie de la reproduction. Mémoire DEA (unpublished). Département de Biologie Animale, Université d'Antananarivo, Madagascar.
- Bauchot, R. (Ed.) (1994). *Snakes: A Natural History*. New York: Sterling Publishing Co., Inc. 220 pp.
- Cadle, J.E. (2003). Colubridae, snakes. In *The Natural History of Madagascar*, pp. 997-1004. Goodman, S.M. & Benstead, J.P. (Eds.). Chicago: University of Chicago Press.
- Crottini, A., Harris, D.J., Irisarri, I.A., Lima, A., Rasamison, S. & Rosa, G.M. (2010). Confirming Domergue: *Ithycyphus oursi* Domergue, 1986 predation upon *Furcifer oustaleti* (Mocquard, 1894). *Herpetology Notes* 3: 127-131
- Crottini, A., Madsen, O., Poux, C., Strauß, A., Vieites, D.R. & Vences, M. (2012) Vertebrate time-tree elucidates the biogeographic pattern of a major biotic change around the K-T boundary in Madagascar. *Proceedings of the National Academy of Sciences* 109: 5358-5363.



Figure 1. Predation of the chameleon *F. major* by a snake, *M. mahfalensis* in Isalo Massif, central-southern Madagascar: **A.** snake seizing the chameleon; **B-C.** snake starting to swallow its prey by the head; **D.** snake unwrapped the chameleon and swallows in gentle motions. Photos by P. Eesebio Bergò.

- Domergue, C.A. (1989). Un serpent venimeux de Madagascar: observation de deux cas de morsure par *Madagascarophis* (Colubridé opisthoglyphe). *Archives de l'Institut Pasteur de Madagascar* 56: 299-311
- Florio, A.M., Ingram C.M., Rakotondravony, H.A., Louis, E.E. & Raxworthy, C.J. (2012). Detecting cryptic speciation in the widespread and morphologically conservative carpet chameleon (*Furcifer lateralis*) of Madagascar. *Journal of Evolutionary Biology* 25: 1399-1414.
- Glaw, F. & Vences, M. (2007). *A Field Guide to the Amphibians and Reptiles of Madagascar*. 3rd edition. Köln: Vences & Glaw. 496 pp.
- Jenkins, R.K.B., Rabearivony, J. & Rakotomanana, H. (2009). Predation on chameleons in Madagascar: a review. *African Journal of Herpetology* 58: 131-136.
- Laduke, T.C., Krause, D.W., Scanlon, J.D. & Kley, N.J. (2010). A Late Cretaceous (Maastrichtian) snake assemblage from the Maevarano Formation, Mahajanga Basin, Madagascar. *Journal of Vertebrate Paleontology* 30: 109-138.
- Mercurio, V., Mattioli, F., Razafindrabe, T.J. & Andreone, F. (2006). A possible attempt of predation of *Liophidium vaillanti* upon *Dromicodryas bernieri* observed in central-southern Madagascar (Serpentes: Colubridae). *Salamandra* 42: 181-183.
- Raxworthy, C.J., Martinez-Meyer, E., Horning, N., Nussbaum, R.A., Schneider, G.E., Ortega-Huerta, M.A. & Peterson, A.T. (2003). Predicting distributions of known and unknown reptile species in Madagascar. *Nature* 426: 837-841.
- Rosa, G.M., Boistel, R., Campantico, E., Gillet, B., Eusebio Bergò, P. & Andreone, F. (2014). Case solved: presence of toxin-secreting oral glands in the lamprophiid snake *Mimophis mahfalensis* (Grandidier, 1867) from Madagascar. *Zoomorphology* 133: 417-423.

Accepted: 25 August 2016