

HERPETOLOGICAL NOTES FROM MAINLAND AND INSULAR GREECE

V. Pérez Mellado *, E.D. Valakos #, M.J. Gil *, F. Guerrero *, J. Lulch \$,
P. Navarro \$ and P. Maragou #

* *Department of Animal Biology, Universidad de Salamanca,
37071- Salamanca. (Spain), e-mail: valentin@gugu.usal.es*

*Department of Biology, Section of Animal & Human Physiology,
University of Athens. GR-15784 Illisia, Athens, Hellas (Greece),
e-mail: evalakos@biology.db.uoa.gr*

\$ *Department of Animal Biology (Parasitology), Universidad de Valencia,
Burjassot, Valencia (Spain)*

INTRODUCTION

Although the herpetofauna of Greece is well known (see for example, Werner, 1938; Wettstein, 1953, 1957; Ondrias, 1968; Chondropoulos, 1986, 1989), several distributional questions still remain unclear, mainly for the mainland but also for islands.

In this paper we present some chorological information from a field trip of one month (April 1990) to mainland and insular Greece (see Figs 1 and 2 and appendix I for a complete list of localities).

RESULTS AND DISCUSSION

Amphibians

Salamandra salamandra. 07.04.90: 1 specimen from Loc. 4 and 1 specimen from Loc. 7 on 09.04.90.

Triturus alpestris. 07.04.90: 1 adult specimen from Loc. 4. First record for the Katara region. Observed on a grassland area at 1705 m.a.s.l.

Bufo bufo. 07.04.90: Several clutches at Loc. 3; 09.04.90: 42 specimens collected at Loc. 6; 09.04.90: 6 specimens from Loc. 7, and 2 specimens from Loc. 9; 10.04.90: 2 specimens from Loc. 10, and 7 specimens from Loc. 13. Also observed at Loc. 8.

Bufo viridis. 13.04.90: 4 specimens from Loc. 16; 18.04.90: 2 specimens collected at Loc. 39, and choruses from Loc. 40 and 41. Also observed at Loc. 18 on 13.04.90.

Rana eppeirotica. 08.04.90: 19 specimens and 09.04.90: 2 specimens from Loc. 5, and 4 specimens from Loc. 6.

The Lake frog present at Lake Ioannina was recorded by Tunner & Heppich (1982) as the "Corfú taxon". However, on Corfú island both species, *Rana eppeirotica* and *Rana ridibunda* coexist (Keymar, 1985, '986). Thus, *Rana eppeirotica* is a cryptic species formerly taken as *Rana ridibunda* (Keymar, 1986). For this reason, the assignment of our sample from Ioannina to this species seems to be difficult without an electrophoretic confirmation. In addition, a morphometrical comparison between these marsh frogs and those from other Greek locations could be helpful in this respect.

Rana graeca. 07.04.90: 7 specimens from Loc. 4; 09.04.90: 4 specimens from Loc. 6; 10.04.90: 1 specimen from Loc. 13. Also observed at Loc. 7 and Loc. 10.

Rana ridibunda; 07.04.90: 14 specimens from Loc. 13, and 1 specimen from Loc. 14; 11.04.90: choruses from Loc. 16; 15.04.90: 1 specimen observed at Loc. 24 and 25 respectively, 7 specimens collected at Loc. 29, 5 specimens collected at Loc. 30, and 2 specimens from Loc. 31; 18.04.90: 1 specimen collected at Loc. 39; 19.04.90: 12 specimens collected at Loc. 33, and 1 specimen from Loc. 36; 20.04.90: 9 specimens from Loc. 37; 26.04.90: 17 specimens collected at Loc. 17, also observed at Loc. 41.

Hyla arborea; 10.04.90: 39 specimens from Loc. 12, and 4 specimens from Loc. 13; 11.04.90: 4 specimens from Loc. 14; 11.04.90: choruses of several individuals at Loc. 15 and 16; 15.04.90: 4 specimens from Loc. 30; 18.04.90: choruses at Loc. 39 and 41; 26.04.90: choruses at Loc. 17.

Reptiles

Emys orbicularis. 14.04.90: 1 specimen observed at Loc. 19; 15.04.90: 1 specimen observed at Loc. 30 (second record for Lesvos Island, the first was from Broggi, 1978); 26.04.90: 2 individuals observed at Loc. 17.

Mauremys caspica. 14.04.90: 1 specimen observed at Loc. 19, and 1 specimen observed at Loc. 20; 15.04.90: 1 specimen observed at Loc. 24; 19.04.90: 6 specimens observed at Loc. 34, and 1 specimen observed at Loc. 38; 20.04.90: several specimens observed, in both cases at Loc. 39, and 2 specimens observed at Loc. 37; 23.04.90: 1 specimen observed at Loc. 43; 24.04.90: 1 specimen from Loc. 45; 26.04.90: extremely common at Loc. 17.

Testudo hermanni. 07.04.90: 1 specimen observed at Loc. 3; 10.04.90: 2 adult specimens observed at Loc. 11.

Testudo marginata. 07.04.90: 1 adult male observed at Loc. 2.

Only recently recorded for some northern localities of mainland and insular Greece (Keymar & Weissinger, 1987).

Anguis fragilis. 08.04.90: 1 adult specimen from Loc. 6.

Ophisaurus apodus. 26.04.90: 1 specimen collected at Loc. 17.

Ablepharus kitaibelli. 06.04.90: 1 specimen from Loc. 1; 24.04.90: 1 specimen from Loc. 44.

Chalcides ocellatus. 19.04.90: 18 specimens from Loc. 36; 20.04.90: 3 specimens from Loc. 33, 1 specimen observed at Loc. 39, and 1 specimen observed from Loc. 42 and also observed at Loc. 38.

Algyroides nigropunctatus. 08.04.90: Two specimens (ad. and subad.) in Loc. 6 and also observed at Loc. 7.

Ophisops elegans ehrenbergii. 14.04.90: 14 specimens from Loc. 19; 14.04.90: 1 specimen from Loc. 18, and 1 specimen from Loc. 20; 15.04.90: 1 specimen observed at Loc. 21, 18 specimens collected at Loc. 23, and several specimens observed at Loc. 25 and 26.

Very common. Recorded from at least in 12 aegean islands (Kühnelt, 1986).

Lacerta trilineata. 14.04.90: 1 specimen observed at Loc. 19; 14.04.90: 2 specimens collected at Loc. 18; 15.04.90: 2 specimens observed at Loc. 21 and 24; 19.04.90: 1 specimen observed at Loc. 34, 1 specimen observed at Loc. 35, 1 specimen observed at Loc. 36, and 1 from Loc. 38; 20.04.90: 2 specimens collected at Loc. 31, several specimens observed at Loc. 39, and 2 specimens collected at Loc. 42; 23.04.90: 1 specimen observed at Loc. 48; 24.04.90: 1 specimen from Loc. 49; 26.04.90: 2 specimens from Loc. 52 and 1 from Loc. 53.

Lacerta vidiris. 07.04.90: 4 specimens collected from Loc. 2; 1 specimen from Loc. 3; 08.04.90: 1 specimen from Loc. 5. and also observed at Loc. 7. 14.04.90: One adult specimen of *Lacerta*, was captured at Loc. 22 in area of Lesvos island with a quite wet environment. Its back pattern, coloration and pholidosis seems to match the species *L. viridis*, but, obviously, we can say nothing conclusive without a biochemical confirmation of its specific status (Mayer & Tiedemann 1981).

Podarcis erhardii. 21.04.90: 46 specimens collected at Loc. 50; 22.04.90: 9 specimens collected at Loc. 51; 26.04.90: 6 individuals observed at Loc. 52 and 53, both from Sifnos island where *Podarcis erhardii* seems to be extremely scarce. Sifnos island is occupied by *P.e.erhardii* (Grillitsch & Tiedemann, 1984). The subspecies *P.e.myconensis* was recorded from Thira by Frör & Beutler (1978).

Podarcis muralis. 08.04.90: 39 specimens from Loc. 5.

Podarcis taurica. 08.04.90. 7 specimens from Loc. 6.

Laudakia stellio. 15.04.90: 1 specimen observed at Loc. 24, and 1 specimen in Loc. 25, 27 and 28 respectively. It was recorded at Lesvos Island by Werner (1938), Ondrias (1968) and Broggi (1978), but its presence was not mentioned by Xyda (1986) who considered that lizards from Samos, Hios and other neighbouring islands have to be included in a new subspecies not nominated in his work.

Cyrtodactylus kotschyi. 23.04.90 and 24.04.90: several specimens observed at Loc. 43, 44, 46, 52 and 53. The species is recorded for the first time by Tiedemann & Häüpl (1982) from Thira (Santorini) and Thirassia.

Hemidactylus turcicus. 20.04.90: 1 specimen collected at Loc. 37.

Tarentola mauritanica. 18.04.90: 1 specimen observed at Loc. 32; 19.04.90: 1 specimen collected at Loc. 34.

Typhlops vermicularis. 10.04.90: 2 specimens from Loc. 11. It is considered as a common species in mainland Greece (Dimitropoulos, 1986).

Coluber gemonensis. 06.04.90: 1 subadult individual from Loc. 1; 20.04.90: 1 specimen collected at Loc. 33. A relatively invariable species of Colubridae, common in mainland Greece (Dimitropoulos, 1986).

Natrix natrix. 08.04.90: 1 specimen from Loc. 5 and 1 specimen from Loc. 6.

Natrix tessellata. 08.04.90: 2 specimens collected at Loc. 5; 09.04.90: 1 specimen from Loc. 10; 10.04.90: 1 specimen from Loc. 13; 26.04.90: 2 individuals observed at Loc. 17.

Vipera lebetina schweizeri. 23.04.90: Observed at Loc. 44. Recorded in the Milos archipelago (Kühnelt, 1986) where it seems to be extremely common but threatened by the persecution from scientific collectors (Dimitropoulos, 1986).

Appendix I. Localities cited in the text (see Figures 1 and 2)

Mainland (Figure 1)

1. Kaza 38° 25' N, 23° 30' E
2. Meteora 39° 50' N, 21° 20' E
3. Meteora (Pinios river)
39° 40' N, 21° 20' E
4. Katara 39° 50' N, 21° 10' E
5. Ioannina (Lake Pamvotida)
38° 25' N, 23° 30' E
6. Ioannina (Lake Pamvotida near to
the islet) 39° 40' N, 20° 40' E
7. Kipi (Zagori) 39° 50' N, 20° 40' E
8. Kopani (Louros river)
39° 20' N, 20° 40' E
9. Fillipiada 39° 20' N, 20° 40' E
10. Trichonida lake 38° 40' N, 21° 20' E
11. Pleuron 38° 20' N, 21° 20' E
12. Souli 38° 00' N, 22° 30' E
13. Stimfalos 37° 50' N, 22° 30' E
14. Kastania 37° 40' N, 22° 20' E
15. Argos 37° 40' N, 22° 40' E
16. Road Argos-Isthmia
37° 40' N, 22° 50' E
17. Shinias (Marathonas)
38° 10' N, 24° 00' E

Lesvos Island (A, Figure 2)

18. Moria 39° 20' N, 26° 30' E
19. Moria 39° 15' N, 26° 30' E
20. Alykes Kallonis (Arisvi)
39° 10' N, 26° 20' E
21. Arisvi 39° 10' N, 26° 15' E
22. Road Mithimna-Vafios
39° 50' N, 26° 10' E
23. Mistegna 39° 20' N, 26° 20' E
24. Mantamados 39° 30' N, 26° 20' E
25. Road Pelopi-Ypsilometopo
39° 20' N, 26° 20' E
26. Road Rodotichos-Kastreli
39° 10' N, 26° 10' E
27. Agra 39° 10' N, 26° 00' E
28. North of Asropotamos river
39° 15' N, 26° 30' E
29. Mylopotamos river
39° 20' N, 26° 20' E
30. Moni Myrtiotissas-Filia
39° 20' N, 26° 10' E
31. Moria 39° 10' N, 26° 25' E

Crete Island (E, Figure 2)

32. Knossos 35° 20' N, 25° 50' E
33. Fodele 35° 20' N, 24° 50' E
34. Road Rizinia-Festos
35° 10' N, 24° 50' E
35. Pano Moulia 35° 00' N, 24° 50' E
36. Gortina 35° 00' N, 24° 50' E
37. Vori 35° 50' N, 24° 40' E
38. Kamilari 35° 00' N, 24° 40' E
39. Lavris (Geropotamos river)
35° 20' N, 24° 50' E
40. Moyri 35° 15' N, 24° 10' E
41. Dramia 35° 20' N, 24° 30' E
42. Rethimno 35° 20' N, 24° 30' E

Milos Island (B, Figure 2)



43. Adamas 36° 40' N, 24° 20' E
44. Filakopi 36° 40' N, 24° 20' E
45. Alykes 36° 40' N, 24° 20' E
46. Provatas 36° 35' N, 24° 20' E
47. Sakoula 36° 35' N, 24° 20' E
48. Sakoula 36° 35' N, 24° 20' E
49. Kimolos (Kimolos Island)
36° 45' N, 24° 20' E

Santorini Island (C, Figure 2)

50. Thira 36° 25' N, 25° 20' E
51. Nea Kamenni 36° 25' N, 25° 15' E

Sifnos Island (D, Figure 2)

52. Apollonia 36° 35' N, 24° 40' E
53. Kamares 37° 00' N, 24° 30' E



REFERENCES

- Broggi, M. (1978). Herpetologische Beobachtungen auf der Inseln Lesbos (Griechenland). *Salamandra* **14**(4): 161-171.
- Chondropoulos, B.P. (1986). A checklist of the Greek reptiles. I. The lizards. *Amphibia-Reptilia* **7**: 217-235.
- Chondropoulos, B.P. (1989). A checklist of the Greek reptiles. II. The snakes. *Herpetozoa* **2** (1/2): 3-36.
- Dimitropoulos, A. (1986). Some notes on the colour and pattern variation of the Greek snake fauna in relation to geographic distribution. *Biologia Gallo-hellenica* **12**: 463-471.
- Frör, E. & Beutler, A. (1978). The herpetofauna of the oceanic island in the Santorini-archipelago, Greece. *Spixiana* **1**: 301-308.
- Grillitsch, H. and Tiedemann (1984). zur Herpetofauna der griechischen Inseln Kea, Spanopoula, Kithnos, Sifnos, Kitriani Cycladen), Alonnisos und Piperi (Nördliche Sporaden). *Ann. Naturhist. Mus. Wien* **86**: 7-28.
- Keymar, P.F. (1985). Additional remarks on: "Some amphibians and reptiles of Corfu with a special note on the occurrence of the nose-horned viper (*Vipera ammodytes*)" (Stafford P. Herptile 9(3) Sep. 1984 pp. 105-108). *Herptile* **10**(1): 25.
- Keymar, P.F. (1986). Die Amphibien und reptilien der Ionischen Region (Griechenland) – Analyse ihrer rezenten Verbreitungsmuster und Überlegungen zu ihrer Ausbreitungsgechichte. *ÖGH-Nachrichten* 1986 (**6/7**): 3-26.
- Keymar, P.F. and Weissingar, H. (1987). Distribution, morphological variation and status of *Testudo marginata* in Greece. In: *Proc. Fourth Ord. Gen. Meet. S.E.H.* J.J. van Gelder, H. Strijbosch & P.J.M. Bergers(eds.) Nijmegen 1987 pp.219-222.
- Kühnelt, W. (1986). Contributions to the knowledge of historical biogeography of Balkan peninsula, especially of Greece. *Biologia Gallo-hellenica* **12**: 71-84.
- Mayer, W. and Tiedemann, F. (1982). Chemotaxonomical investigation in the collective genus *Lacerta* (Lacertidae, Sauria) by means of protein electroforesis. *Amphibia-Reptilia* **2**: 346-356.
- Ondrias, J. C. (1968). Liste des Amphibiens et des Reptiles de Grèce. *Biologia Gallo-hellenica*. **1**: 111-135.
- Tiedemann, F. and Häupl, M. (1982). *Cyrtodactylus kotschy* (Steindachner, 1870) in the Santorin archipelago. *Amphibia-Reptilia* **3**: 377-381.
- Tunner, H.G. and Heppich, S. (1982). A genetic analysis of water frogs from Greece: evidence for the existence of a cryptic species. *Z.zool. Syst. Evolut.-forsch.* **20**: 209-233.
- Werner, F. (1938). Die Amphibien und Reptilien Griechenlands. *Zoologica* **35**: 1-117.
- Wettstein, O. (1953). Herpetologia aegaea. *Sitz.-ber. Östeer. Akad. Wiss., Math.-naturwiss. Kl.* **166**: 123-164.
- Xyda, A. (1986). Supplementary evidence on the biometry and ecology of the lizard *Stellio stellio* of Greece and Cyprus. *Biologia Gallo-hellenica* **12**: 451-458.