

THE HERPETOLOGICAL JOURNAL

Volume 16, Number 2 2006

CONTENTS

Full Papers

- | | | |
|--|---|---------|
| A modelling approach to determine a translocation scenario for the endangered New Zealand frog <i>Leiopelma hamiltoni</i> | M. D. TOCHER, D. FLETCHER & P. J. BISHOP | 97–106 |
| Variation between populations in the diet of the Mediterranean lizard <i>Lacerta perspicillata</i> | A. PERERA, V. PÉREZ-MELLADO, M. A. CARRETERO & D. J. HARRIS | 107–113 |
| Anuran temporal occupancy of a temporary pond in the Atlantic rain forest, south-eastern Brazil | P. A. ABRUNHOSA, H. WOGEL & J. P. POMBAL JR. | 115–122 |
| Long distance movements by <i>Caiman crocodilus yacare</i> : implications for management of the species in the Brazilian Pantanal | Z. CAMPOS, M. COUTINHO, G. MOURÃO, P. BAYLISS & W. E. MAGNUSSON | 123–132 |
| Age, body size and growth of <i>Lacerta agilis boemica</i> and <i>L. strigata</i> : a comparative study of two closely related lizard species based on skeletochronology | E. S. ROITBERG & E. M. SMIRINA | 133–148 |
| Ecomorphological guilds in anuran larvae: an application of geometric morphometric methods | M. F. VERA CANDIOTI | 149–162 |
| Ecological functions of the foam nests of the Japanese treefrog, <i>Rhacophorus arboreus</i> (Amphibia, Rhacophoridae) | T. KUSANO, A. SAKAI & S. HATANAKA | 163–169 |
| Life in the water: ecology of the Jacarerana lizard, <i>Crocodylus amazonicus</i> | M. MARTINS | 171–176 |
| Reproductive ecology of a Mediterranean ratsnake, the ladder snake <i>Rhinechis scalaris</i> (Schinz, 1822) | J. M. PLEGUEZUELOS & M. FERICHE | 177–182 |
| A new species of arboreal <i>Leptopelis</i> (Anura: Hyperoliidae) from the forests of western Kenya | J. KÖHLER, B. A. BWONG, S. SCHICK, M. VEITH & STEFAN LÖTTERS | 183–189 |
| Systematics of the nose-horned viper (<i>Vipera ammodytes</i> , Linnaeus, 1758) | L. TOMOVIC | 191–201 |
| The tadpole of <i>Physalaemus fernandezae</i> (Anura: Leptodactylidae) | L. ALCALDE, G. S. NATALE & R. CAJADE | 203–211 |
| Geographic variation and taxonomic status of the southernmost populations of <i>Liophis miliaris</i> (Linnaeus, 1758) (Serpentes: Colubridae) | A. R. GIRAUDO, V. ARZAMENDIA & P. CACCIALI | 213–220 |
| Genetic evidence for two distinct species within the Italian endemic <i>Salamandrina terdigitata</i> (Bonnaterre, 1789) (Amphibia: Urodela: Salamandridae) | D. CANESTRELLI, F. ZANGARI & G. NASCETTI | 221–227 |
| <i>Short Note</i> | | |
| First description of the acoustic repertoire of <i>Rana italica</i> (Anura, Ranidae) | E. RAZZETTI, R. SACCHI & J. E. PLATZ | 229–231 |