

Infection by the chytrid fungus *Batrachochytrium dendrobatidis* in the yellow belly frog (*Elachistocleis bicolor*) from Argentina

María Luz Arellano¹, Daiana P. Ferraro^{2,3}, Mónica M. Steciow¹ & Esteban O. Lavilla⁴

¹Instituto de Botánica Carlos Spegazzini, Buenos Aires, Argentina

²Museo de La Plata, Buenos Aires, Argentina

³Museo Argentino de Ciencias Naturales "Bernardino Rivadavia", Buenos Aires, Argentina

⁴Instituto de Herpetología, Fundación Miguel Lillo, Tucumán, Argentina

Infection by the chytrid fungus *Batrachochytrium dendrobatidis* was detected in a dead adult of the yellow belly frog (*Elachistocleis bicolor*) in Misiones province, Argentina. The analysis of the skin through standard histological procedures revealed the presence of chytrid sporangia in different stages. This is the first report of *B. dendrobatidis* occurring in hot and humid lowlands in northeastern Argentinian Atlantic forest, and the first case of infection for the frog genus *Elachistocleis*. The importance of this finding is discussed in light of the particular conservation importance of the eco-region involved, and the coinciding emergence of locally introduced American bullfrogs.

Key words: Anura, amphibians, disease, Microhylidae, Misiones

Chytridiomycosis is considered an emerging disease (Daszak et al., 2003) linked to mortalities of wild amphibian populations in many areas including Europe, South America, Central America, North America, New Zealand, Australia and Africa (Bishop, 2000; Bosch et al., 2001; Bradley et al., 2002; Weldon & du Preez, 2004; Barrionuevo & Mangione, 2006; Lips et al., 2006). This disease is caused by the chytrid fungus *Batrachochytrium dendrobatidis* (*Bd*, Longcore et al., 1999; Phylum Chytridiomycota, Order Rhizophydiales; Letcher et al., 2006). At present, *B. dendrobatidis* has been reported in eight South American countries, including Argentina, Bolivia, Brazil, Colombia, Ecuador, Peru, Uruguay and Venezuela (Ron & Merino, 2000; Bonaccorso et al., 2003; Mazzoni et al., 2003; Carnaval et al., 2005; Herrera et al., 2005; Lampo et al., 2006; Seimon et al., 2005; Barrionuevo et al., 2008; Ruiz & Rueda-Almonacid, 2008; Velásquez et al., 2008). In Argentina,

infections by *Bd* have been recorded in five anuran species: *Atelognathus patagonicus*, *Leptodactylus gracilis*, *L. ocellatus*, *Telmatobius atacamensis* and *T. pisanoi* (Herrera et al., 2005; Barrionuevo & Mangione, 2006; Fox et al., 2006; Ghirardi et al., 2009), taxa associated with high altitudes and/or latitudes or with low temperatures.

In this contribution we report the presence of the fungus on an individual of *Elachistocleis bicolor* from Misiones Province (Argentina). *Elachistocleis bicolor* (Guerín-Méneville, 1838), is a small, ovoid-shaped microhylid, widely distributed in Uruguay, northeastern Argentina, southern Brazil and Paraguay, and southwestern Bolivia (IUCN 2008), and is considered "Least Concern" by Lavilla et al. (2000) and IUCN (2008). The province of Misiones includes all the Atlantic forest remnants in Argentina (Giraudó, 2005). Although it covers only 1.1 % of the total area of the country, it is one of the richest in biodiversity in Argentina, and one of the most threatened on a global scale (Giraudó et al., 2005).

On 26 November 2005 one individual of the yellow belly frog was found dead (without signs of have been killed by vehicle traffic) on the margin of a temporal pond on National Route N° 12, on the road to Posadas (27°26'373"S, 56°01'43"W, 122 m a.s.l.), Misiones province, Argentina (Figure 1). The specimen was fixed in 10% formalin and then maintained in 70% ethanol. The ventral pelvic and ventral hind limb skin was dehydrated, paraffin-embedded and cut (Microm GmbH HM 400 R) into five-micron-thick sections for histological analysis. Skin cuts were stained with haematoxylin and eosin and were examined with a light microscope (Olympus BX 40 Optical Co. Ltd, Tokyo, Japan) We followed the guidelines described in Berger et al. (1999) to diagnose infection produced by *Bd*. The voucher specimen is housed in the herpetological collection of the Museo de La Plata, Buenos Aires, Argentina (MLP A. 4631).

Histological examination of the epidermis of the yellow belly frog showed evidence of fungal infection. Spherical and ovoid zoosporangia, empty or containing zoospores, were identified in the keratinized cell layer of the epidermis (Fig. 2). Many of the zoosporangia, ranging from 5 to 10 µm in diameter, showed the discharge tube that releases zoospores. The infected areas included mild to moderate hyperkeratosis and areas of focal erosion adjacent to the infection. These observations agree with *Bd* infection as described by Berger et al. (1999). The stage of the infection was identified as "light", containing clusters of 4–6 spherical sporangia (Herrera et al., 2005).

Although the study is based on one specimen, the presence of *B. dendrobatidis* in *E. bicolor* from Misiones (Argentina) is relevant for several reasons. It represents the southernmost record of *Bd* in the Paranean forest ecoregion (*sensu* Burkart et al., 1999). The Atlantic forest, considered one of the five biodiversity hotspots originally reported for South America (Myers et al., 2000; Conservation International, 2005), is currently reduced to 7–8% of its original size (Galindo Leal & Gusmão Câmara, 2005), and not only houses the highest diversity of am-



Fig. 1. Report site (indicated by star) of the yellow belly frog infected by *Bd* in Misiones province, northeastern Argentina. The triangles represent four *Lithobates catesbeianus* (American bullfrog) localities (Pereyra et al., 2006).

phibians in Argentina and Paraguay, but also is one of the richest batrachological areas in Brazil (Brusquetti & Lavilla, 2006; Cochran, 1955; Cei, 1980). The Atlantic forest is also characterized by its high level of endemic species (Plací & Di Bitetti, 2005). While only about 3% and 10% of the original cover exist in Brazil and Paraguay, respectively, Argentina still maintains 50% of the original forest (Holz & Placci, 2005; Giraudo et al., 2005). Previously, *Bd* was reported from several localities from Jaqueira (Pernambuco, Brazil; Carnaval et al., 2005, 2006; Toledo et al., 2006) at a straight-line distance of about 2900 km to Misiones.

Our study area has a humid-subtropical climate (*sensu* Köppen, 1936), where the mean minimum temperature is 21.8 °C for the hottest month (January, local summer) and 11.1 °C for the coldest (June, local winter; statistics for the period 1981–1991; Servicio Meteorológico Nacional, 2008). Our finding constitutes the first record of *Bd* in a lowland and subtropical area of Argentina during spring. Previous records of *Bd* in the country were associated with colder temperatures, either due to winter records (Buenos Aires Province, 11 m a.s.l., 4 July 2002; Herrera et al., 2005), mid or high altitudes (Salta Province, 3885 m a.s.l., Tucumán Province, 2352 m a.s.l., Barrionuevo & Magione, 2006; Córdoba Province 800 m a.s.l., Ghirardi et al., 2009), or high latitudes (Neuquén Province, Laguna Blanca, 39°S, 70°W, 1276 m a.s.l., Fox et al., 2006). It is interesting to note that reports of *Bd* around the world are scarce in lowlands (Mazzoni et al., 2003; Carnaval et al.,

2006; Puschendorf et al., 2006) and mainly occur in farmed animals and/or under low temperatures.

Interestingly, the presence of *Bd* in the Paranean forest of Misiones coincides with the finding in the same ecoregion of feral populations of the American bullfrog *Lithobates catesbeianus* (Pereyra et al., 2006), which can act as an efficient pathogen carrier (Kats & Ferrer, 2003; Garner et al., 2006). *Lithobates catesbeianus* is considered resistant to the chytrid fungus (Daszak et al., 2004; Hanselmann et al., 2004), and infected specimens have been recorded in three South American countries, including Brazil (Garner et al., 2006), Uruguay (Mazzoni et al., 2003) and Venezuela (Hanselmann et al., 2004). Aside from Misiones, populations of bullfrogs in Argentina have also been recorded in San Juan (Sanabria et al., 2005), Córdoba (Akmentins et al., 2009), Buenos Aires (Barraso et al., 2009) and Salta (Akmentins & Cardozo, 2009), with *Bd* reports in the latter three provinces (Barrionuevo & Mangione, 2006; Ghirardi et al., 2009; Herrera et al., 2005). Specific studies of *Bd* infection in *L. catesbeianus* are necessary to confirm if this anuran species is acting as a carrier of the disease in Misiones province.

Up to now, no other *Bd* infections have been recorded in Misiones province. At the pond where the specimen was collected, a further 16 anuran species (adults and/or tadpoles) have been recorded (Diego Baldo, pers. comm.). Studies on species susceptibility to *Bd* are needed to quantify its potential to affect amphibian populations in Argentina, considering that this fungal disease has been suggested to cause death in whole frog communities from Australian and Central American Forests (Berger et al., 1998; Lips, 1999).

Acknowledgements. We thank D. Baldo, F. Kolenc, C. Borteiro, M. Tedros and E. Kubisch for their help during field trips and D. Baldo for providing valuable information about species in the study area. We wish to thank P. Grilli for his support during manuscript development. Fauna authorities of Misiones province provided collection permit. DPF acknowledges ANPCyT (Argentina) by grant PICT06-223.




Fig. 2. Histological section of ventral skin of the yellow belly frog. a) Mature zoosporangium (mz) containing zoospores in the stratum corneum. b) Mature zoosporangium almost empty; cs: stratum corneum; gs: stratum granulosum. Scale bar (a, b) = 18 μ m.

REFERENCES

- Akmentins, M.S. & Cardoso, D.E. (2009). American bullfrog *Lithobates catesbeianus* (Shaw, 1802) invasion in Argentina. *Biological Invasions*. DOI: 10.1007/s10530-009-9515-3.
- Akmentins, M.S., Pereyra, L.C. & Lescano, J.N. (2009). Primer registro de una población asilvestrada de rana toro (*Lithobates catesbeianus*) en la provincia de Córdoba, Argentina. Notas sobre la biología de la especie. *Cuadernos de Herpetología* 23, 25–32.
- Barraso, D.A., Cajade, R., Nenda, S.J., Baloriani, G. & Herrera, R. (2009). Introduction of the American bullfrog *Lithobates catesbeianus* (Anura Ranidae) in natural and modified environments: an increasing conservation problem in Argentina. *South American Journal of Herpetology* 4, 69–75.
- Barrionuevo, J.S., Aguayo, R. & Lavilla, E.O. (2008). First record of chytridiomycosis in Bolivia (*Rhinella quechua*; Anura: Bufonidae). *Diseases of Aquatic Organisms* 82, 161–163.
- Barrionuevo, S. & Mangione, S. (2006). Chytridiomycosis in two species of *Telmatobius* (Anura: Leptodactylidae) from Argentina. *Diseases of Aquatic Organisms* 73, 171–174.
- Berger, L., Speare, R., Daszak, P., Green, D.E., Cunningham, A.A., Gonnin, C.L., Slocombe, R., Ragan, M.A., Hyatt, A.D., McDonald, K.R., Hines, H.B., Lips, K.R., Marantelli, G. & Parkes, H. (1998). Chytridiomycosis causes amphibian mortality associated with population declines in the rain forests of Australia and Central America. *Proceedings of the National Academy of Sciences* 95, 9031–9036.
- Berger, L., Speare, R. & Kent, A. (1999). Diagnosis of chytridiomycosis in amphibians by histologic examination. www.jcu.edu.au/school/phtm/PHTM/frogs/histo/chhisto.htm (accessed 20 November 2008).
- Bishop, P. (2000). Chytrid fungi identified from dying frogs in New Zealand. *Froglog* 39, 1–2.
- Bonaccorso, E., Guayasamin, J.M., Méndez, D. & Speare, R. (2003). Chytridiomycosis as a possible cause of population declines in *Atelopus cruciger* (Anura: Bufonidae). *Herpetological Review* 34, 331–334.
- Bosch, J.I., Martínez-Solano, I. & García-Paris, M. (2001). Evidence of a chytrid fungus infection involved in the decline of the midwife toad (*Alytes obstetricans*) in protected areas of Central Spain. *Biological Conservation* 97, 331–337.
- Bradley, G.A., Rosen, P.C., Sredl, J.M. & Jones, T.R. (2002). Chytridiomycosis in native Arizona frogs. *Journal of Wildlife Diseases* 38, 206–212.
- Brusquetti, F. & Lavilla, E.O. (2006). Lista comentada de los anfibios de Paraguay. *Cuadernos de Herpetología* 20 (2), 3–79.
- Burkart, R., Bárbaro, N., Sánchez, R.O. & Gómez, D.A. (1999). *Eco-Regiones de la Argentina*. Buenos Aires: Administración de Parques Nacionales.
- Carnaval, A.C.O.Q., Puschendorf, R., Peixoto, O.L., Verdade, V.K. & Rodrigues, M.T. (2006). Amphibian chytrid fungus broadly distributed in the Brazilian Atlantic rain forest. *EcoHealth* 3, 41–48.
- Carnaval, A.C.O.Q., Toledo, L.F., Haddad, C.F.B. & Britto, F.B. (2005). Chytrid fungus infects high-altitude stream-dwelling *Hylodes magalhaesi* (Leptodactylidae) in the Brazilian Atlantic rainforest. *Froglog* 70, 3–4.
- Cei, J.M. (1980). Amphibians of Argentina. *Monitore Zoologico Italiano (N.S.)*, Monogr. 2, 1–609.
- Cochran, D.M. (1955). Frogs of southeastern Brazil. *United States National Museum Bulletin* 206, 1–423.
- Conservation International (2005). *Biodiversity Hotspots*. Available at <http://www.biodiversityhotspots.org/xp/Hotspots> (accessed 11 August 2009).
- Daszak, P., Cunningham, A.A. & Hyatt, A.D. (2003). Infectious disease and amphibian population declines. *Diversity and Distributions* 9, 141–150.
- Daszak, P., Strieby, A., Cunningham, A.A., Longcore, J.E., Brown, C.C. & Porter, D. (2004). Experimental evidence that the bullfrog (*Rana catesbeiana*) is a potential carrier of chytridiomycosis, an emerging fungal disease of amphibians. *Herpetological Journal* 14, 201–207.
- Fox, S.F., Greer, A.L., Torres-Cervantes, R. & Collins, J.P. (2006). First case of ranavirus-associated morbidity and mortality in natural populations of the South American frog *Atelognathus patagonicus*. *Diseases of Aquatic Organisms* 72, 87–92.
- Galindo Leal, C. & Gusmão Câmara, I. (2005). El estado del “Hotspot” Bosque Atlántico: una visión general. In *El Bosque Atlántico en Paraguay, Biodiversidad, Amenazas y Perspectivas*, 3–12. Cartes, J.L. (ed.). Asunción: Conservation International – Center for Applied Biodiversity Science – Guyra Paraguay.
- Garner, T.W.J., Perkins, M.W., Govindarajulu, P., Seglie, D., Walker, S., Cunningham, A.A. & Fisher, M.C. (2006). The emerging amphibian pathogen *Batrachochytrium dendrobatidis* globally infects introduced populations of the North American bullfrog, *Rana catesbeiana*. *Biology Letters* 2, 455–459.
- Ghirardi, R., Lescano, J.N., Longo, M.S., Robledo, G., Steciow, M.M. & Perotti, M.G. (2009). *Batrachochytrium dendrobatidis* in Argentina: first record in *Leptodactylus gracilis* and another record in *Leptodactylus ocellatus*. *Herpetological Review* 40, 175–176.
- Giraud, A. (2005). Dinâmica da perda da biodiversidade na Mata Atlântica argentina: uma introdução. In *Mata Atlântica, Biodiversidade, Ameaças e Perspectivas*, 137–140. Galindo Leal, C. & Gusmão Câmara, I (eds). Belo Horizonte: Conservação Internacional.
- Giraud, A.R., Povedano, H., Belgrano, M.J., Krauczuk, E., Pardiñas, U., Miquelarena, A., Ligier, D., Baldo, D. & Castelino, M. (2005). Status da biodiversidade da Mata Atlântica de interior da Argentina. In *Mata Atlântica, biodiversidade, ameaças e perspectivas*, 160–180. Galindo Leal, C. & Gusmão Câmara, I (eds). Belo Horizonte: Conservação Internacional.
- Hanselmann, R., Rodríguez, A., Lampo, M., Fajardo-Ramos, L., Aguirre, A.A., Kilpatrick, A.M., Rodríguez, J.P. & Daszak, P. (2004). Presence of an emerging pathogen of amphibians in introduced bullfrogs (*Rana catesbeiana*) in Venezuela. *Biological Conservation* 120, 115–119.
- Herrera, R.A., Steciow, M.M. & Natale, G.S. (2005).

- Chytrid fungus parasitizing the wild amphibian *Leptodactylus ocellatus* (Anura: Leptodactylidae) in Argentina. *Diseases of Aquatic Organisms* 64, 247–252.
- Holz, S. & Placci, G. (2005). Raízes socioeconômicas da perda da biodiversidade em Misiones. In *Mata Atlântica, Biodiversidade, Ameaças e Perspectivas*, 207–226. Galindo Leal, C. & Gusmão Câmara, I (eds). Belo Horizonte: Conservação Internacional.
- IUCN (2008). *2008 IUCN Red List of Threatened Species*. www.iucnredlist.org (accessed 18 February 2009).
- Kats, L.B. & Ferrer, R.P. (2003). Alien predators and amphibian declines: review of two decades of science and the transition to conservation. *Diversity and Distributions* 9, 99–110.
- Köppen, W. (1936). Das geographische System der Klimate. In *Handbuch der Klimatologie*, 1–44. Köppen, W. & Geiger, G. (eds). Berlin: Borntraeger.
- Lampo, M., Barrio-Amorós, C. & Han, B. (2006). *Batrachochytrium dendrobatidis* infection in the recently rediscovered *Ateopus mucubajiensis* (Anura, Bufonidae), a critically endangered frog from the Venezuelan Andes. *EcoHealth* 3, 299–302.
- Lavilla, E.O., Ponssa, M.L., Baldo, D., Basso, N., Bosso, A., Céspedes, J., Chebez, J.C., Faivovich, J., Ferrari, L., Lajmanovich, R., Langone, J.A., Peltzer, P., Úbeda, C., Vaira, M. & Vera Candioti, F. (2000). Categorización de los anfibios de Argentina. In *Categorización de los Anfibios y Reptiles de la República Argentina*, 11–34. Lavilla, E.O., Richard, E. & Scrocchi, G.J. (eds). Tucumán: Asociación Herpetológica Argentina.
- Letcher, P.M., Powell, M.J., Churchill, P.F. & Chambers, J.G. (2006). Ultrastructural and molecular phylogenetic delineation of a new order, the Rhizophydiales (Chytridiomycota). *Mycological Research* 110, 898–915.
- Lips, K.R. (1999). Mass mortality of the anuran fauna at an upland site in Panama. *Conservation Biology* 13, 117–125.
- Lips, K.R., Burrowes, P.A., Mendelson, J.R. & Parra-Olea, G. (2006). Amphibian declines in Latin America: widespread population declines, extinctions, and impacts. *Biotropica* 37, 163–165.
- Longcore, J.C., Pessier, A.P. & Nichols, D.K. (1999). *Batrachochytrium dendrobatidis* gen. et sp. nov., a chytrid pathogenic to amphibians. *Mycologia* 91, 219–227.
- Mazzoni, R., Cunningham, A.A., Daszak, P., Apolo, A., Perdomo, E. & Speranza, G. (2003). Emerging pathogen of amphibians in frogs (*Rana catesbeiana*) farmed for international trade. *Emerging Infectious Diseases* 9, 995–998.
- Myers, N., Mittermeier, R.A., Mittermeier, C.G., da Fonseca, G.A.B. & Kent, J. (2000). Biodiversity hotspots for conservation priorities. *Nature* 403, 853–858.
- Pereyra, M.O., Baldo, D. & Kraukzuc, E.R. (2006). La “rana toro” en la selva atlántica interior Argentina: un nuevo problema de conservación. *Cuadernos de Herpetología* 20, 37–40.
- Plací, G. & Di Bitetti, M. (2005). Situación ambiental en la ecoregión del Bosque Atlántico del Alto Paraná (Selva Paranaense). In *La Situación Ambiental Argentina 2005*, 195–225. Brown, A., Martínez Ortiz, U., Acerbi, M. & Corcuera, J. (eds). Buenos Aires: Fundación Vida Silvestre Argentina.
- Puschendorf, R., Bolaños, F. & Chaves, G. (2006). The amphibian chytrid fungus along an altitudinal transect before the first reported declines in Costa Rica. *Biological Conservation* 132, 136–142.
- Ron, S.R. & Merino, A. (2000). Declinación de anfibios del Ecuador: información general y primer reporte de chytridiomycosis para Sudamérica. *Froglog* 42, 2.
- Ruiz, A. & Rueda-Almonacid, V.J. (2008). *Batrachochytrium dendrobatidis* and chytridiomycosis in anuran amphibians of Colombia. *EcoHealth* 5, 27–33.
- Sanabria, E.A., Quiroga, L.B. & Acosta, J.C. (2005). Introducción de *Rana catesbeiana* Shaw (rana toro), en ambientes pre-cordilleranos de la provincia de San Juan, Argentina. *Multiequina* 14, 67–70.
- Seimon, T.A., Hoernig, G., Sowell, P., Halloy, S. & Seimon, A. (2005). Identification of chytridiomycosis in *Telmatobius marmoratus* at 4,450 m in the Cordillera Vilcanota of southern Peru. In *Studies on the Andean Frogs of the Genera Telmatobius and Batrachophrynus*, 275–283. Lavilla, E.O. & de la Riva, I. (eds). Monografías de Herpetología 7. Valencia: Asociación Herpetológica Española.
- Servicio Meteorológico Nacional (2008). Argentina: Misiones: Posadas. www.smn.gov.ar/?mod=clima&id=30&provincia=Misiones&ciudad=Posadas (accessed 2 May 2008).
- Toledo, L.F., Britto, F.B., Araújo, O.G.S., Giasson, L.M.O. & Haddad, C.F.B. (2006). The occurrence of *Batrachochytrium dendrobatidis* in Brazil and the inclusion of 17 new cases of infection. *South American Journal of Herpetology* 1, 185–191.
- Velásquez-E., B.E., Castro, F., Bolívar-G., W. & Herrera, M.I. (2008). Infección por el hongo quitridio *Batrachochytrium dendrobatidis* en anuros de la cordillera occidental de Colombia. *Herpetotrópicos* 4, 65–70.
- Weldon, C.H. & du Preez, L.H. (2004). Decline of the Kihansi spray toad, *Nectophrynoides asperginis*, from the Udzungwa Mountains, Tanzania. *Froglog* 62, 2–3.

Accepted: 11 September 2009