

HABITAT SELECTION BY THE LIZARD *LACERTA LEPIDA* IN A MEDITERRANEAN OAK FOREST

AURORA M. CASTILLA^{1,3} AND DIRK BAUWENS²

¹ Museo Nacional de Ciencias Naturales, c/J. Gutiérrez Abascal 2, E-28006 Madrid, Spain

² Institute of Nature Conservation, Kiewitdreef 3, B-3500 Hasselt, Belgium

³ present address: Dept. Biology, University of Antwerp (U.I.A.), Universiteitsplein 1, B-2610 Wilrijk, Belgium

(Accepted 21.8.90.)

ABSTRACT

We studied habitat selection by *Lacerta lepida* in an open, degraded Mediterranean *Quercus ilex*-forest using two different methods. First, the frequency of lizard observations in distinct habitat types was compared with habitat availability. Although lizards were seen in a wide range of habitats, a preference for sites with a complex vertical vegetation structure was evident. We also quantified structural features of the habitat at sighting spots of adult lizards. Both univariate and multivariate analyses indicated that lizards preferred sites with a relatively high coverage of high (>2 m) vegetation, but with some low (<50 cm) vegetation, and where some rocks were present.

INTRODUCTION

Lacerta lepida, the largest European lacertid lizard (adult SVL: 140-230 mm), is distributed throughout most of Spain and Portugal, southern France and extreme north-west Italy (Bischoff *et al.*, 1984). It is a diurnal, heliothermic, ground-dwelling lizard that forages actively on a large variety of mainly arthropod prey (Castilla *et al.*, 1991). This lizard occurs from sea-level up to ca. 2000 m (Seva, 1982; Filella, 1983), and is found in a wide variety of both uncultivated and man-made habitats, including coastal sand-plains and dunes, arable land, olive groves, orchards, vineyards, grassland, shrubland, etc. (Arnold & Burton, 1976; Seva, 1982; Bischoff *et al.*, 1984; Barbadillo, 1987). Due to its seemingly ubiquitous presence in different habitats, it is generally considered as not exhibiting strong habitat preferences. However, quantitative data on habitat use are very scarce. Only Busack & Visnaw (1989) provide quantitative information on the presence of *L. lepida* in different habitats within the province of Cádiz (southern Spain). These authors concluded that lizards inhabited different habitat types in direct proportion to their availability (but see Discussion).

We here report on a study of the habitat selection patterns of *L. lepida* in an open Mediterranean oak woodland in central Spain. Very high densities are usually attained in this macrohabitat, which is often composed of distinct habitat types. We combine two complementary approaches to the study of habitat preference. First, we examine lizard sighting frequency in the different habitat categories in relation to their availability. Second, we attempt to identify the cues that guide habitat occupation through a comparison of quantitative features of the habitat structure between sites occupied by lizards and available habitats.

MATERIALS AND METHODS

The study was conducted between March and September 1985-1986 in an area of degraded oak woodland ("encinar adehesado") near Navas del Rey (40°23'N, 4°15'W, province of Madrid, Spain). The study site consists of several well-recognisable habitats, shaped mainly through varying degrees of past and current human exploitation (sheep and cattle grazing, burning).

HABITAT CATEGORIES

Observations on habitat utilisation were made while one of us walked along a fixed 3700 m transect, which crossed all habitat categories. The route was walked 125 times between 07.00-22.00 hr (Mean European Time), covering the entire daily activity period of lizards. Upon sighting of a lizard, we recorded: date, time, size (Age) class [juvenile: < 70 mm snout-vent length (SVL); subadult: 71-139 mm SVL; adult: SVL ≥ 140 mm], and habitat category.

We established 12 representative habitat categories: 1) Dry, homogeneous grassland; 2) low (<50 cm), mixed shrubland with young oak (*Quercus ilex*), dwarf-shrubs and grasses; 3) dry areas with a patchy covering of dwarf-shrubs (mainly *Lavandula stoechas*); 4) moist areas with abundant cover of rushes (*Juncus sp.*); 5) sites with a patchy covering of broom bushes (*Lygos sphaerocarpa*); 6) dense bushes of brambles (*Rubus sp.*); 7) almond (*Prunus dulcis*) orchard; 8) riverside areas with ashes (*Fraxinus angustifolia*); 9) riverside areas with willows (*Salix atrocinerea*); 10) isolated *Juniperus oxycedrus* trees with dense grassy undergrowth; 11) open, degraded oak forest (*Quercus ilex*) with grassy undergrowth; 12) open evergreen forest of mixed composition (*Quercus ilex*, *Juniperus oxycedrus*) and with a heterogeneous undergrowth of shrubs and grasses.

We estimated the availability of the distinct habitat categories by recording their presence at 25 m intervals along the transect line. These frequency counts were converted to the expected number of lizard observations under the null-hypothesis that sighting frequency in each habitat was directly proportional to its availability.

SELECTION OF HABITAT FEATURES

We quantified structural features of the habitat at sighting spots of 66 adult lizards. Habitat availability was estimated through sampling 51 sites that were situated at 70 m intervals along the transect line.

For each habitat sample we located 5 points: the spot where the lizard was first sighted, and the endpoints of two orthogonal 6 m lines that intersected at the sighting spot. The direction of these lines was determined haphazardly by throwing a pen on the ground. Each of these points served as the centre of a circle with 1 m radius. Within each circle we estimated the following 11 variables: 1-3) percentage cover at ground level of rock/stone, sand and litter; 4-8) percentage cover of the vegetation layers with heights <25 cm, 25-50 cm, 50-100 cm, 100-200 cm and >200cm; 9) maximum height of the vegetation; 10) distance towards the nearest rock and 11) distance towards the nearest patch of vegetation that would potentially provide shelter to predators. Three additional variables were calculated; 12) total vegetation cover: summation of the vegetation cover over the 5 layers; 13) vertical diversity: calculated over the 5 vegetation layers with the Shannon-Wiener formula; 14) horizontal diversity: calculated over the habitat variables that cover the ground (rock/stone, sand, litter, vegetation <25 cm) with the Shannon-Wiener formula.

We calculated the mean value for each variable in the 5 sampling points. This value estimates the average habitat structure within a circular area with 4 m radius, surrounding the spot where the lizard was seen. Our data hence provide a description of habitat structure for an area, rather than for a particular point, that is a portion of the lizard's home range.

We performed a principal component analysis on the correlation matrix of the habitat variables to reduce the dimensionality of the habitat space. We use *F*-tests, and *t*-tests (equal variances) or Mann-Whitney *U*-tests (unequal variances) to evaluate the statistical significance of differences in variances and means among lizard observation sites and availability samples, both for the original variables and the scores on the principal components.

RESULTS

HABITAT CATEGORIES

Lizard size classes did not differ in sighting frequency in the distinct habitat categories ($X^2 = 14.99$, 22 df, $P > 0.80$). We did not detect seasonal differences in habitat occupation ($X^2 = 29.07$, 33 df, $P > 0.10$), and therefore lumped all data for further analysis.

Lizards were observed in all habitats. Sighting frequency in the distinct habitat categories was not directly proportional to their availability ($X^2 = 327.97$, 11 df, $P < 0.001$; Fig. 1). We saw more lizards than expected in the mixed forest, in mixed shrubland, and near bramble bushes. Lizards seemed to avoid grasslands and, to a lesser extent, vegetations dominated by rushes, patches of broom bushes, and oak forests (Fig. 1).




Fig. 1. Observed ($N = 509$) and expected sighting frequencies of *L. lepida* in distinct habitat categories.

Gr: grassland; Sh: mixed shrubland; Lv: vegetations with *Lavandula stoechas*; Ru: moist areas with rushes; Ls: sites with *Lygos sphaerocarpa*; Br: dense bushes of brambles; Al: almond orchard; Fx: riverside areas with ashes; Sx: riverside areas with willows; Ju: isolated *Juniperus oxycedrus*-trees with dense grassy undergrowth; Qu: open oak forest; Mx: mixed evergreen forest.

SELECTION OF HABITAT FEATURES

Structural features of sites occupied by adult lizards and of available microhabitats are summarised in Table 1. Note that for many variables the observation sites had a higher variance, but a similar mean value than the availability samples. This indicates that lizards were seen more often than expected at sites with low values, and at places with high scores for these habitat characteristics.

	Observed	Available	P var	P means
% cover sand	21.1±15.2	27.5±15.4	NS	NS
rock	21.1±16.8	15.5±12.6	**	NS
litter	8.7±15.1	5.8±11.1	**	NS
% cover veg. <25cm	49.0±21.3	50.9±17.2	*	NS
25-50cm	29.1±18.1	30.8±13.8	*	NS
50-100cm	13.1±11.1	11.8±8.9	*	NS
100-200cm	7.4±6.6	7.1±7.9	NS	NS
>200cm	16.2±21.6	8.5±9.7	***	NS
tot. veg. cover	114.9±43.4	109.2±39.4	NS	NS
max. veg. height (cm)	246.2±171.5	158.3±90.0	***	**
distance veg. (cm)	146.1±94.8	249.0±172.4	***	***
distance rock (cm)	238.3±250.4	329.2±241.9	NS	*
vertical div.	0.84±0.22	0.79±0.21	NS	NS
horizontal div.	0.59±0.18	0.61±0.14	*	NS
PC1	-0.09±2.44	0.12±1.75	**	NS
PC2	0.48±1.93	-0.63±1.64	NS	**
PC3	0.19±1.27	-0.25±1.19	NS	=0.06

TABLE 1. Summary (mean ± 1 SD) of structural habitat features at lizard observation sites and availability samples, and probability of differences between their variances (*F*-test) and means (*t*-test or Mann-Whitney *U*-test). N = 66 for lizard observation sites; N = 51 for availability samples. *: *P*<0.05; **: *P*<0.01; ***: *P*>0.001; NS: *P*>0.05.

A principal component analysis was used to obtain a more integrated picture of the habitat attributes. The first three principal components accounted respectively for 31, 24 and 10% of the total variance of the raw data. The first principal component (PC1) was strongly positively correlated with percent cover of the vegetation layers <25 cm, 25-50 cm and 50-100 cm, total vegetation cover, vertical diversity, and distance to the nearest rock.

	PC1	PC2	PC3
% cover sand	-0.052	-0.667	-0.314
rock	-0.566	0.303	0.605
litter	-0.514	0.627	-0.333
% cover veg. <25cm	0.845	-0.135	0.013
25-50cm	0.896	0.105	0.013
50-100cm	0.785	0.323	0.100
100-200cm	0.332	0.489	-0.084
>200cm	-0.357	0.795	-0.250
tot. veg. cover	0.844	0.480	-0.085
max. veg. height (cm)	-0.331	0.821	-0.280
distance veg. (cm)	-0.185	-0.489	-0.481
distance rock (cm)	0.507	-0.403	-0.438
vertical div.	0.599	0.550	-0.185
horizontal div.	-0.465	0.001	-0.412

TABLE 2. Correlations between habitat characteristics and the first three principal components. Only correlation coefficients significant at $\alpha = 0.001$ and higher than 0.50 were considered in the interpretation of the principal axes, and are shown in boldface.

It was negatively correlated with percent cover by stones/rocks and by litter at ground level (Table 2). This represents a gradient from habitats with extensive cover by stones/rocks and litter, and sparse vegetation near the ground towards sites with abundant vegetation at <100 cm above ground level. The second principal axis was positively correlated with percent cover by vegetation >200 cm high, maximum vegetation height, vertical diversity and percent cover of litter, and exhibited a strong negative correlation with percent cover by sand at ground level. Thus, PC2 describes a gradient from open, sandy sites to locations with upright bushes or trees. The third principal component showed a strong positive correlation with percent cover by rocks (Table 2).

We examined habitat occupation by considering the projections (= scores) of the different samples on the component axes. The range of the scores for lizard sighting spots was similar to that of the available sites on the three first principal components. Comparison of the distribution of scores on PC1 reveals a higher variance, but a similar average, for lizard sighting spots than for available sites (Table 1). This indicates that more lizards than expected were seen at both extremes of the gradient represented by PC1, i.e. at sites with few low vegetation and many rocks on the one hand, and at places with much low vegetation on the other hand. Scores on PC2 were significantly higher for observation sites (Table 1), indicating that lizards preferred habitats with upright trees or bushes, and that they avoided open, sandy areas. Lizard sighting spots had a marginally non-significant higher average score on PC3, suggesting a tendency for lizards to select sites with much cover by rocks at ground level.

DISCUSSION

Our data, collected within a restricted area of degraded Mediterranean woodland, indicate that *L. lepida* used the entire range of available habitats. Nevertheless, lizards exhibited a preference for certain habitat types. They were observed more often than expected in the mixed forest, in mixed shrubland and near bramble bushes, and they avoided grasslands. Busack & Visnaw (1989) reported data on habitat use collected in a large geographical area (province of Cádiz, Spain), and distinguished between several broad habitat categories, based on their type of land use. *L. lepida* was found in most of these biotopes and it was concluded that it did not exhibit any habitat preference (Busack & Visnaw, 1989). Reanalysis of their data, using an appropriate frequency-dependent statistical procedure, indicates that lizards do not inhabit the different habitat types proportionally to their abundance ($X^2 = 76.32$, 12 df. *P* <0.001). More lizards than expected were seen in sclerophyllous shrub and grassland, whereas sighting frequency was lower than expected in agriculturally unproductive areas [see Table 4 in Busack & Visnaw (1989)]. Hence, although *L. lepida* should be considered as a habitat generalist, its preference for certain biotopes is evident both on a local and a regional scale.

Our study of the quantitative features of habitat structure indicates that lizards were seen throughout the range of available sites. This again shows that lizards

occupied all structural habitats that were available in the study area. However, sighting frequency was higher than expected at sites with upright (>2 m) trees and bushes, at places with many stones or rocks and at habitats with much low (<1 m) vegetation. Lizards tended to avoid open, sandy areas with little vegetation. These results point to a preference for sites with a rather complex habitat structure, although lizards are definitely not restricted to such places.

Several factors might invoke a preference for habitats with a high structural diversity. The presence of trees or bushes creates a sun-shade gradient. This should facilitate the shuttling between sunlit and shaded sites, a conspicuous aspect of the thermoregulatory behaviour of Lacertid lizards (Avery, 1976; Arnold, 1987; Van Damme *et al.*, 1987). Rocks and/or stones might also serve a thermoregulatory function. On repeated occasions we observed lizards basking on rocks or stones near abundant low vegetation. The variable orientation of stony surfaces should facilitate postural adjustments that increase net radiation intake, and a close contact with their sun-warmed substrates enhances heat gain through conduction. At the same time, rock-crevices and holes between piled stones provide shelter to predators. In addition, it might be hypothesised that the number of invertebrate prey might be highest in structurally complex habitats (Strijbosch, 1988).

REFERENCES

- Arnold, E. N. (1987). Resource partitioning among lacertid lizards in southern Europe. *J. Zool. (Lond.) (B)*, **1**, 739-782.
- Arnold, E. N. and Burton, J. A. (1978). *A field guide to the reptiles and amphibians of Britain and Europe*. Collins, London.
- Avery, R.A. (1976). Thermoregulation, metabolism and social behaviour in Lacertidae. In: *Morphology and biology of reptiles*. Bellairs, A.d'A. and Cox, C.B. (Eds.). Linnean Society Symposium Ser. no. 3, London, pp. 245-259.
- Barbadillo, L. J. (1987). *La guía de Incafo de los anfibios y reptiles de la Península Ibérica, Islas Baleares y Canarias*. Incafo, Madrid.
- Bischoff, W., Cheylan, M. and Böhme, W. (1984). *Lacerta lepida* Daudin 1802 — Perleidechse. In: *Handbuch der Reptilien und Amphibien Europas. Band 2/1*. Böhme, W. (Eds.). AULA-Verlag, Wiesbaden, pp 181-210.
- Busack, S. D. and Visnaw, J. (1989). Observations on the natural history of *Lacerta lepida* in Cádiz Province, Spain. *Amphibia-Reptilia*, **10**, 201-213.
- Castilla, A. M. (1989). *Autoecología del Lagarto ocelado (Lacerta lepida)*. Ph. D. Thesis. Universidad Autónoma de Madrid, Madrid.
- Castilla, A. M., Bauwens, D. and Llorente, G. A. (1991). Diet composition of the lizard *Lacerta lepida* in central Spain. *J. Herpet.* **25**, 30-36.
- Christian, K., Porter, W. P. and Tracy, C. R. (1986). Core areas within the home ranges of Galapagos land iguanas. *Conolophus pallidus*. *J. Herpet.*, **20**, 272-276.
- Christian, K. A. and Waldschmidt, S. (1984). The relationship between lizard home range and body size: a reanalysis of the data. *Herpetologica*, **40**, 68-75.
- Filella, E. (1983). Nota sobre *Lacerta lepida lepida*. Daudin 1802 (Sauria, Lacertidae). *Bull. Soc. Ictio. Herp.* **5**, 18-19.
- Grant, B. W. and Dunham, A. E. (1988). Thermally imposed time constraints on the activity of the desert lizard *Sceloporus merriami*. *Ecology*, **69**, 167-176.
- Huey, R. B. and Pianka, E. R. (1977). Seasonal variation in thermoregulatory behaviour and body temperature of diurnal Kalahari lizards. *Ecology*, **58**, 1066-1075.
- Seva, E. (1982). *Taxocenosis de Lacertidos en un arenal costero alicantino*. Publ. Universidad de Alicante, Alicante.
- Stamps, J. A. (1977). Social behaviour and spacing patterns in lizards. In: *Biology of the reptilia*. Vol 7. Gans, C. and Tinkle, D. W. (Eds.). Academic Press, London, pp 265-334.
- Strijbosch, H. (1988). Habitat selection of *Lacerta vivipara* in a lowland environment. *Herpet. J.*, **1**, 207-210.
- Van Damme, R., Bauwens, D. and Verheyen, R. F. (1987). Thermoregulatory responses to environmental seasonality by the lizard *Lacerta vivipara*. *Herpetologica*, **43**, 405-415.

ACKNOWLEDGEMENTS

This study was supported by grants of the I.N.I.A. and the "Comunidad Autónoma de Madrid" (to AMC). Preparation of the manuscript was aided by a postdoctoral grant of the C.S.I.C. (to AMC), and by logistic support of IBERIA, National Spanish Airlines.