



## Detection of *Salmonella enterica* in a sand lizard (*Lacerta agilis*, Linnaeus, 1758) city population

Krzysztof Dudek<sup>1</sup>, Ryszard Koczura<sup>2</sup>, Monika Dudek<sup>3</sup>, Zofia Anna Sajkowska<sup>4</sup>  
& Anna Ekner-Grzyb<sup>5</sup>

<sup>1</sup>Department of Zoology, Institute of Zoology, Poznań University of Life Sciences, Wojska Polskiego 71 C, 60-625 Poznań, Poland

<sup>2</sup>Department of Microbiology, Faculty of Biology, Adam Mickiewicz University in Poznań, Umultowska 89, 61-614 Poznań, Poland

<sup>3</sup>Laboratory of Neurobiology, Institute of Zoology, Poznań University of Life Sciences, Wojska Polskiego 71 C, 60-625 Poznań, Poland

<sup>4</sup>Laboratory of Biological and Natural Education, Faculty of Biology, Adam Mickiewicz University in Poznań, Umultowska 89, 61-614 Poznań, Poland

<sup>5</sup>Department of Behavioural Ecology, Faculty of Biology, Adam Mickiewicz University in Poznań, Umultowska 89, 61-614 Poznań, Poland

Salmonellosis is one of the most urgent public health problems across the world. Reptiles are a known reservoir of *Salmonella* spp. and in some regions they are also associated with human salmonellosis. This concerns especially popular pet reptiles, e.g., turtles or bearded dragons; however, there is also a need for studies regarding wild reptiles as a pathogen source. In this study, sand lizards (*Lacerta agilis*) were investigated as a potential reservoir of *Salmonella* spp. in Poznań, Poland, using cloacal swabs and faecal samples. Moreover, clonal analysis of the isolates was conducted using ERIC-PCR fingerprinting. Thirty eight lizards were investigated, nine of which (24%) proved positive for *S. enterica* subsp. *houtenae*. The prevalence level was lower than previously observed in exotic species (up to above 40%). Two clones were present in several lizards. Specimens with similar clones were captured at the same location and time, suggesting horizontal transfer of bacterial strains between lizards. Because the isolated subspecies of *Salmonella* is very rarely reported as a causative agent of human salmonellosis, sand lizards seem to pose little or no threat for public health.

**Key words:** clonal analysis, environmental microbiology, infectious disease, reptile-associated salmonellosis

*Salmonella* spp. is a regular component of the microbial flora of the digestive tract in reptiles (Mitchell & Shane, 2001, see also Benskin et al., 2009 for birds). In mammals their presence leads to salmonellosis (Buxton, 1957) and reptiles have therefore been linked to salmonellosis outbreaks in humans (Warwick et al., 2001; Hassl & Benyr, 2003; Mermin et al., 2004; Bauwens et al., 2006; Bertrand et al., 2008). *Salmonella enterica* subsp. *enterica* is the most pathogenic subspecies, and transmission from pet reptiles to humans has previously been reported (Woodward et al., 1997; Hidalgo-Vila et al., 2007; Pedersen et al., 2009; Chen et al., 2010).

Prevalence of *Salmonella* spp. in reptiles is often high (e.g., Geue & Löschner, 2002: 54.1%; Briones et al., 2004: 41.5%), and transovarial transfer may occur from mother to clutch (Chiodini, 1982). Other studies however also report that *Salmonella* spp. is absent (Geue & Löschner, 2002).

Due to their close contact with humans, studies which link human salmonellosis with reptiles have previously large focused on pet reptiles (e.g., *Trachemys scripta*; Warwick et al., 2001; Nagano et al., 2006; CDC, 2008), and our knowledge on the occurrence of *Salmonella* spp. in wild European reptiles is still very incomplete (e.g., *Emys orbicularis*: Hidalgo-Vila et al., 2007, *Natrix natrix*: Wuthe et al., 1979; Rostami et al., 2009, *Vipera berus*: Wuthe et al., 1979). The sand lizard (*Lacerta agilis*) is the most widespread reptile species in Europe (Bischoff, 1988), but our knowledge about the presence of *Salmonella* spp. is restricted to a single individual investigated which was tested positively for *Salmonella sofia* (currently *S. enterica* subsp. *salamae*) several decades ago (Koopman & Janssen, 1973). The aim of this study was to document the occurrence of *Salmonella* in a Polish population of *L. agilis*.

The study was carried out in May–June 2014 in two localities in Poznań, Poland (near Rusałka Lake, 52°25'38.86N, 16°52'14.63E and on the edge of Morasko forest, 52°28'03.03N, 16°55'50.54E) in habitats which are strongly effected by anthropogenic activities. *Lacerta agilis* is a short-legged, ground-dwelling diurnal lizard of around 66–73mm snout-vent-length (SVL, Ekner et al., 2008).

Lizards were captured using nets or by hand. SVL of caught lizards were measured to classify them as adult (>45 mm SVL), sub-adults (35–45 mm SVL), or juveniles (<35 mm, Gvozdik, 2000; Dudek et al., 2015). Juveniles were lizards in their first year of life, subadults were in their second year of life, and adults were at least three years old. Individuals were sexed based on the presence

Correspondence: Krzysztof Dudek (dudekk@gmail.com)

Dice (Opt:1.00%) (Tol 1.0%-1.0%) (H&gt;0.0% S&gt;0.0%) [0.0%-100.0%]




Fig. 1. Dendrogram showing genetic relatedness of bacterial isolates determined by ERIC-PCR typing. Cluster 1 comprises isolates J10 and J11; cluster 2 comprises isolates J33 and J36.

of femoral pores and the expanded gonadal area in the tail base in males. During the mating season males also have green body colouration.

After taking cloacal swabs, all lizards were placed overnight in separate sterile boxes. On the next day, the faecal samples were collected using Amies Agar Gel transport swabs (Oxoid) and processed within four hours. To avoid pseudoreplication, the lizards were marked using medical cautery units (following Ekner et al., 2011), and each individual was used only once. After sample collection, all individuals were released at the place of their capture. Cloacal swabs and stool specimens were inoculated into Rappaport-Vassiliadis Enrichment Broth (Oxoid) and incubated 48 hours at 42°C. If bacterial growth was observed, the culture was inoculated onto Brilliant Green Agar (Oxoid) and incubated 18–24 hours at 35°C. Lactose-negative isolates were identified using API 20E kits (bioMérieux) and complementary tests for utilisation of malonate, mucate and salicine (Popoff & Le Minor, 2005).

Isolates were genotyped using ERIC-PCRs, with primers complementary to enterobacterial repetitive intergenic consensus sequences. Genomic DNA was isolated using the Genomic Mini Kit (A&A Biotechnology). PCR reactions with primers ERIC1R and ERIC 2 were conducted in a C1000 thermal cycler (Bio-Rad) following Versalovic et al. (1991). Amplicons were separated on 2% agarose gels, and banding patterns were analysed using GelCompar II 3.5 software (Applied Maths) using the Dice Similarity Coefficient and UPGMA clustering. Isolates sharing DNA fingerprinting patterns above 95% similarity were considered clones. The experiments were done in duplicate. Prevalence and confidence limits (95% CI) for binary, presence-absence, data were calculated in Microsoft Excel 2013 (Microsoft).

A total of 38 individuals (13 females, 20 males, and 5 juveniles) were captured. Specimens taken from 10 lizards yielded bacterial growth in Rappaport-Vassiliadis Enrichment Broth. Nine of these cultures grew as typical red-coloured lactose-negative colonies on Brilliant Green

Agar, and were identified with API 20E as *S. enterica*. Complementary tests revealed that all nine isolates were malonate- and mucate-negative and salicine-positive, which allowed identifying them as *S. enterica* subsp. *houtenae*. Prevalence was estimated as 0.237 (95% CI=0.102–372).

The ERIC fingerprints consisted of 8–18 bands ranging from 180 bp to 6300 bp in size. The dendrogram showed the presence of two clusters with 100% similarity (Fig. 1). The first cluster comprised isolates no. J10 and J11, cultured from lizards collected on 04 July 2014; the second one consisted of isolates no. J33 and J36, cultured from lizards collected on 10 July 2014. With less than 90% similarity, the remaining five isolates were genetically unrelated.

Nine out of 38 *L. agilis* individuals were positive for *S. enterica* in faecal samples. Such a prevalence rate is below most previously reported values (e.g., Geue & Löschner, 2002 on captive individuals: 47.4% for lizards, distributed across Agamidae 60%, Chamaeleonidae 71.4%, Iguanidae 62.1%, Phrynosomatidae and Scinidae 33.3%, Gekkonidae 16.7%, Crotophytidae and Pochrotidae 0%; Colubridae 96.2%, Boidae 56.1%). All salmonellae isolates cultured from stool and cloacal swabs were identified as *S. enterica* subsp. *houtenae*. Strains of this subspecies are mostly isolated from cold-blooded animals and their environment (Geue & Löschner, 2002; Popoff & Le Minor, 2005; Bauwens et al., 2006; Bertrand et al., 2008; Pedersen et al., 2008; Hydeskov et al., 2013; Gay et al., 2014), and have also been found in intestines of wild boars (Chiari et al., 2013; Zottola et al., 2013) and birds (Millán et al., 2004). Overall in reptiles, *S. enterica* subsp. *houtenae* is the third most commonly found *Salmonella* taxon after *S. enterica* subsp. *enterica* and *S. enterica* subsp. *Diarizonae*, ahead of *S. enterica* subsp. *salamae*, *S. enterica* subsp. *arizonae*, and *S. bongori* (Bertrand et al., 2008).

Clonal analysis showed the presence of two clusters consisting of isolates with identical ERIC fingerprintings, i.e. clones. These clones were cultured from lizards

captured at the same day and location (Morasko and Rusałka Lake, respectively). This suggests that strains of *Salmonella* spp. can be transmitted horizontally between lizards. There is a need to investigate whether transovarial transfer of *Salmonella* spp. can occur in sand lizards, as is the case for snakes (Chiodini, 1982).

The frequent occurrence of *S. enterica* subsp. *houtenae* likely poses little or no threat for public health, as this subspecies is only very rarely reported in pet reptiles and as a causative agent of human salmonellosis (Bertrand et al., 2008; Hoszowski et al., 2000, 2012; Sadkowska-Todys & Czarkowski, 2013). There is however a need for further studies on other common reptile species as potential reservoirs for *Salmonella*.

**Acknowledgements:** We thank the Editor and two anonymous reviewers for their constructive comments. This work was supported by a grant for young scientists no. 507.511.06 from the Ministry of Science and Higher Education of Poland. Lizards were caught by using a method that meets Polish legal standards and in accordance with the rules set by the Ethics Committee for Animal Experimentation (LKE 25/2013).

## REFERENCES

- Bauwens, L., Vercammen, F., Bertrand, S., Collard, J.M., De Ceuster, S. (2006). Isolation of *Salmonella* from environmental samples collected in the reptile department of Antwerp Zoo using different selective methods. *Journal of Applied Microbiology* 101, 284–289.
- Benskin, C.M., Wilson, K., Jones, K., Hartley, I.R. (2009). Bacterial pathogens in wild birds: a review of the frequency and effects of infection. *Biological Reviews* 84, 349–373.
- Bertrand, S., Rimhanen-Finne, R., Weill, F.X., Rabsch, W., et al. (2008). *Salmonella* infections associated with reptiles: the current situation in Europe. Euro surveillance: bulletin European sur les maladies transmissibles. *European Communicable Disease Bulletin* 13, 717–727.
- Bischoff, W. (1988). Zur Verbreitung und Systematik der Zauneidechse, *Lacerta agilis* Linnaeus, 1758. *Mertensiella* 1, 11–30.
- Briones, V., Téllez, S., Goyache, J., Ballesteros, C., et al. (2004). *Salmonella* diversity associated with wild reptiles and amphibians in Spain. *Environmental Microbiology* 6, 868–871.
- Buxton, A. (1957). *Salmonellosis in Animals. A Review*. Commonwealth Agricultural Bureaux. Farnham Royal, Nr. Slough, Bucks., England.
- Centers for Disease Control and Prevention, (2008). Multistate outbreak of human *Salmonella* infections associated with exposure to turtles - United States, 2007–2008. *MMWR. Morbidity and mortality weekly report* 57, 69.
- Chen, C.Y., Chen, W.C., Chin, S.C., Lai, Y.H., et al. (2010). Prevalence and antimicrobial susceptibility of salmonellae isolates from reptiles in Taiwan. *Journal of Veterinary Diagnostic Investigation* 22, 44–50.
- Chiari, M., Zanoni, M., Tagliabue, S., Lavazza, A., Alborali, L.G. (2013). *Salmonella* serotypes in wild boars (*Sus scrofa*) hunted in northern Italy. *Acta Veterinaria Scandinavica* 55, 42.
- Chiodini, R.J. (1982). Transovarian passage, visceral distribution, and pathogenicity of salmonella in snakes. *Infection and Immunity* 36, 710–713.
- Dudek, K., Sajkowska, Z., Gawałek, M., Ekner-Grzyb, A. (2015). Using body condition index can be an unreliable indicator of fitness: a case of sand lizard *Lacerta agilis* Linnaeus, 1758 (Sauria: Lacertidae). *Turkish Journal of Zoology* 39, 182–184.
- Ekner, A., Sajkowska, Z., Dudek, K., Tryjanowski, P. (2011). Medical cautery units as a permanent and non-invasive method of marking lizards. *Acta Herpetologica* 6, 229–236.
- Ekner, A., Majlath, I., Majlathova, V., Hromada, M., et al. (2008). Densities and morphology of two co-existing lizard species (*Lacerta agilis* and *Zootoca vivipara*) in extensively used farmland in Poland. *Folia Biologica* 56, 165–171.
- Gay, N., Le Hello, S., Weill, F.X., de Thoisy, B., Berger, F. (2014). *Salmonella* serotypes in reptiles and humans, French Guiana. *Veterinary Microbiology* 170, 167–171.
- Geue, L., Löschner, U. (2002). *Salmonella enterica* in reptiles of German and Austrian origin. *Veterinary Microbiology* 84, 79–91.
- Gvozdik, L. (2000). Seasonal activity, sex ratio, and abundance in a population of *Lacerta agilis* Linnaeus, 1758 from the Czech Republic. *Herpetozoa* 13, 165–169.
- Hassl, A., Benyr, G. (2003). Hygienic evaluation of terraria inhabited by amphibians and reptiles: cryptosporidia, free-living amebas, salmonella. *Wiener Klinische Wochenschrift* 115, 68–71.
- Hidalgo-Vila, J., Díaz-Paniagua, C., de Frutos-Escobar, C., Jiménez-Martínez, C., Pérez-Santigosa, N. (2007). *Salmonella* in free living terrestrial and aquatic turtles. *Veterinary Microbiology* 119, 311–315.
- Hoszowski, A., Wasyl, D., Truszczynski, M. (2000). *Salmonella* serovars determined in the National Veterinary Research Institute in 1994–1998. *Bulletin of the Veterinary Institute in Pulawy* 44, 33–38.
- Hoszowski, A., Skarzyńska, M., Wasyl, D., Zajac, M., et al. (2012). Serowary *Salmonella* występujące u zwierząt, w żywności i paszach w Polsce w latach 2005–2010. *Medycyna Weterynaryjna* 68, 411–417.
- Hydeskov, H.B., Guardabassi, L., Aalbæk, B., Olsen, K.E.P., et al. (2013). *Salmonella* prevalence among reptiles in a zoo education setting. *Zoonoses and Public Health* 60, 291–295.
- Koopman, J.P., Janssen, F.G.J. (1973). The occurrence of salmonellas and lactose-negative *Arizonas* in reptiles in The Netherlands, and a comparison of three enrichment methods used in their isolation. *Journal of Hygiene Cambridge* 71, 363–371.
- Mermin, J., Hutwagner, L., Vugia, D., Shallow, S., et al. (2004). Reptiles, amphibians, and human *Salmonella* infection: a population-based, case-control study. *Clinical Infectious Diseases* 38, 253–261.
- Millán, J., Aduriz, G., Moreno, B., Juste, R.A., Barral, M. (2004). *Salmonella* isolates from wild birds and mammals in the Basque Country (Spain). *Revue scientifique et technique International Office of Epizootics* 23, 905–911.
- Mitchell, M.A., Shane, S.M. (2001). *Salmonella* in reptiles. In *Seminars in avian and exotic pet medicine* 10, 55–35.
- Nagano, N., Oana, S., Nagano, Y., Arakawa, Y. (2006). A severe *Salmonella enterica* serotype Paratyphi B infection in a child related to a pet turtle, *Trachemys scripta elegans*. *Japanese*

- Journal of Infectious Diseases* 59, 132.
- Pedersen, K., Lassen-Nielsen, A.M., Nordentoft, S., Hammer, A.S. (2009). Serovars of *Salmonella* from captive reptiles. *Zoonoses Public Health* 56, 238–242.
- Popoff, M.Y., Le Minor, L.E. (2005). Genus *Salmonella*. In: Brenner, D.J., Krieg, N.R., Staley, J.T. (eds.) *Bergey's Manual of Systematic Bacteriology. Volume 2: The Proteobacteria, Part B: The Gammaproteobacteria*. pp. 764–799. Springer-Verlag, New York.
- Rostami, A., Memarian, I., Salehi, T.Z., Dehghan, M.M., et al. (2009). The report of *Salmonella typhimurium* in *Natrix natrix*; detection and identification by culture and multiplex PCR methods. In Proceedings of the International Conference on Diseases of Zoo and Wild Animals, Beekse Bergen, The Netherlands, 20th–24th May 2009. pp. 312–316. Leibniz Institute for Zoo and Wildlife Research.
- Sadkowska-Todys, M., Czarkowski, M.P. (2013). Salmonellosis in Poland in 2011. *Przegląd Epidemiologiczny* 67, 467–472.
- Versalovic, J., Koeuth, T., Lupski, J.R. (1991). Distribution of repetitive DNA sequences in eubacteria and application to fingerprinting of bacterial genomes. *Nucleic Acids Research* 19, 6823–6831.
- Warwick, C., Lambiris, A.J., Westwood, D., Steedman, C. (2001). Reptile-related salmonellosis. *Journal of the Royal Society of Medicine* 94, 124.
- Woodward, D.L., Khakhria, R., Johnson, W.M. (1997). Human salmonellosis associated with exotic pets. *Journal of Clinical Microbiology* 35, 2786–2790.
- Wuthe, H.H., Rohde, R., Aleksić, S., Schubert, C., Wuthe, S. (1979). Zentralblatt für Bakteriologie, Parasitenkunde, Infektionskrankheiten und Hygiene. Erste Abteilung Originale. Reihe A: Medizinische Mikrobiologie und Parasitologie, 243, 412–418.
- Zottola, T., Montagnaro, S., Magnapera, C., Sasso, S., et al. (2013). Prevalence and antimicrobial susceptibility of *Salmonella* in European wild boar (*Sus scrofa*); Latium Region – Italy. *Comparative Immunology, Microbiology and Infectious Disease* 36, 161–168.

Accepted: 23 March 2015