



## **One gecko's pain is another gecko's gain: is the Moorish gecko *Tarentola mauritanica* becoming invasive in France?**

Julien Renet<sup>1\*,\*\*</sup>, Théo Dokhelar<sup>2,\*\*\*</sup> & Nicolas Dubos<sup>3,\*\*\*\*</sup>

<sup>1</sup> Fauna studium, Scientific consulting, 04200 Sisteron, France

<sup>2</sup> CEFE, University of Montpellier, CNRS, EPHE, IRD, 34090 Montpellier, France


<sup>3</sup> The Jacob Blaustein Institutes for Desert Research, Sde-Boqer Campus, Ben Gurion University of the Negev, Midreshet Ben-Gurion 8499000, Israel

\*Corresponding author; e-mail: [julien.renet13@hotmail.fr](mailto:julien.renet13@hotmail.fr)


\*\* ORCID 0000-0002-7975-2963

\*\*\* ORCID 0000-0002-8780-767X

\*\*\*\* ORCID 0000-0002-1224-2598



**Figure S1.** Comparison of the average number of observations of the two species between transects over the 8 sampling years. Boxes indicate the 25%, 50% (median) and 75% quartiles. Lines show minimum and maximum values. Different letters represent significant differences between groups.



**Figure S2.** Comparison of abundance variation between *Euleptes europaea* and *Tarentola mauritanica* for each transect during sampling years. Boxes indicate the 25%, 50% (median) and 75% quartiles. Lines show minimum and maximum values excluding outliers (indicated as dots).